

NEW YORK NON-NATIVE PLANT INVASIVENESS RANKING FORM

Scientific name: Cyperus difformis L. USDA Plants Code: CYDI
 Common names: Variable flatsedge
 Native distribution: Southern Europe, Asia, Africa
 Date assessed: 20 July 2009
 Assessors: Gerry Moore
 Reviewers: LIISMA SRC
 Date Approved: September 9, 2009 Form version date: 10 July 2009

New York Invasiveness Rank: Moderate (Relative Maximum Score 50.00-69.99)

Distribution and Invasiveness Rank (Obtain from PRISM invasiveness ranking form)			
Status of this species in each PRISM:		Current Distribution	PRISM Invasiveness Rank
1	Adirondack Park Invasive Program	Not Assessed	Not Assessed
2	Capital/Mohawk	Not Assessed	Not Assessed
3	Catskill Regional Invasive Species Partnership	Not Assessed	Not Assessed
4	Finger Lakes	Not Assessed	Not Assessed
5	Long Island Invasive Species Management Area	Restricted	Moderate
6	Lower Hudson	Not Assessed	Not Assessed
7	Saint Lawrence/Eastern Lake Ontario	Not Assessed	Not Assessed
8	Western New York	Not Assessed	Not Assessed

Invasiveness Ranking Summary (see details under appropriate sub-section)		Total (Total Answered*) Possible	Total
1	Ecological impact	40 (<u>20</u>)	6
2	Biological characteristic and dispersal ability	25 (<u>25</u>)	18
3	Ecological amplitude and distribution	25 (<u>25</u>)	15
4	Difficulty of control	10 (<u>7</u>)	1
Outcome score		100 (<u>77</u>) ^b	40 ^a
Relative maximum score †			51.95
New York Invasiveness Rank §		Moderate (Relative Maximum Score 50.00-69.99)	

* For questions answered "unknown" do not include point value in "Total Answered Points Possible." If "Total Answered Points Possible" is less than 70.00 points, then the overall invasive rank should be listed as "Unknown."

† Calculated as 100(a/b) to two decimal places.

§ Very High >80.00; High 70.00–80.00; Moderate 50.00–69.99; Low 40.00–49.99; Insignificant <40.00
 Not Assessable: not persistent in NY, or not found outside of cultivation.

A. DISTRIBUTION (KNOWN/POTENTIAL): Summarized from individual PRISM forms

A1.1. Has this species been documented to persist without cultivation in NY? (reliable source; voucher not required)	
<input checked="" type="checkbox"/>	Yes – continue to A1.2
<input type="checkbox"/>	No – continue to A2.1
A1.2. In which PRISMs is it known (see inset map)?	
<input type="checkbox"/>	Adirondack Park Invasive Program
<input type="checkbox"/>	Capital/Mohawk
<input type="checkbox"/>	Catskill Regional Invasive Species Partnership
<input type="checkbox"/>	Finger Lakes
<input checked="" type="checkbox"/>	Long Island Invasive Species Management Area
<input type="checkbox"/>	Lower Hudson
<input type="checkbox"/>	Saint Lawrence/Eastern Lake Ontario
<input type="checkbox"/>	Western New York



NEW YORK NON-NATIVE PLANT INVASIVENESS RANKING FORM

Documentation: Specimen collected in 2002 from Queens at BBG. Other specimens recently collected from New Jersey. Author has seen this species at several sites in New Jersey in recent years. Appears to be rapidly spreading.

Sources of information:

Brooklyn Botanic Garden, 2009; author's pers. obs.

A2.1. What is the likelihood that this species will occur and persist outside of cultivation, given the climate in the following PRISMs? (obtain from PRISM invasiveness ranking form)

Not Assessed	Adirondack Park Invasive Program
Not Assessed	Capital/Mohawk
Not Assessed	Catskill Regional Invasive Species Partnership
Not Assessed	Finger Lakes
Very Likely	Long Island Invasive Species Management Area
Not Assessed	Lower Hudson
Not Assessed	Saint Lawrence/Eastern Lake Ontario
Not Assessed	Western New York

Documentation: Species recently identified in LIISMA PRISM.

Sources of information (e.g.: distribution models, literature, expert opinions):

Brooklyn Botanic Garden, 2009.

If the species does not occur and is not likely to occur in any of the PRISMs, then stop here as there is no need to assess the species. Rank is "Not Assessable."

A2.2. What is the current distribution of the species in each PRISM? (obtain rank from PRISM invasiveness ranking forms)

	Distribution
Adirondack Park Invasive Program	Not Assessed
Capital/Mohawk	Not Assessed
Catskill Regional Invasive Species Partnership	Not Assessed
Finger Lakes	Not Assessed
Long Island Invasive Species Management Area	Restricted
Lower Hudson	Not Assessed
Saint Lawrence/Eastern Lake Ontario	Not Assessed
Western New York	Not Assessed

Documentation:

Sources of information:

Brooklyn Botanic Garden, 2009; Weldy & Werier, 2009.

A2.3. Describe the potential or known suitable habitats within New York. Natural habitats include all habitats not under active human management. Managed habitats are indicated with an asterisk.

<p>Aquatic Habitats</p> <input type="checkbox"/> Salt/brackish waters <input type="checkbox"/> Freshwater tidal <input type="checkbox"/> Rivers/streams <input checked="" type="checkbox"/> Natural lakes and ponds <input checked="" type="checkbox"/> Vernal pools <input checked="" type="checkbox"/> Reservoirs/impoundments*	<p>Wetland Habitats</p> <input checked="" type="checkbox"/> Salt/brackish marshes <input checked="" type="checkbox"/> Freshwater marshes <input type="checkbox"/> Peatlands <input type="checkbox"/> Shrub swamps <input type="checkbox"/> Forested wetlands/riparian <input checked="" type="checkbox"/> Ditches* <input type="checkbox"/> Beaches and/or coastal dunes	<p>Upland Habitats</p> <input type="checkbox"/> Cultivated* <input type="checkbox"/> Grasslands/old fields <input type="checkbox"/> Shrublands <input type="checkbox"/> Forests/woodlands <input type="checkbox"/> Alpine <input checked="" type="checkbox"/> Roadsides*
--	--	---

Other potential or known suitable habitats within New York:

Documentation:

Sources of information:

Brooklyn Botanic Garden, 2009; author's pers. obs.

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

B. INVASIVENESS RANKING

Questions apply to areas similar in climate and habitats to New York unless specified otherwise.

1. ECOLOGICAL IMPACT

1.1. Impact on Natural Ecosystem Processes and System-Wide Parameters (e.g. fire regime, geomorphological changes (erosion, sedimentation rates), hydrologic regime, nutrient and mineral dynamics, light availability, salinity, pH)

- A. No perceivable impact on ecosystem processes based on research studies, or the absence of impact information if a species is widespread (>10 occurrences in minimally managed areas), has been well-studied (>10 reports/publications), and has been present in the northeast for >100 years. 0
- B. Influences ecosystem processes to a minor degree (e.g., has a perceivable but mild influence on soil nutrient availability) 3
- C. Significant alteration of ecosystem processes (e.g., increases sedimentation rates along streams or coastlines, reduces open water that are important to waterfowl) 7
- D. Major, possibly irreversible, alteration or disruption of ecosystem processes (e.g., the species alters geomorphology and/or hydrology, affects fire frequency, alters soil pH, or fixes substantial levels of nitrogen in the soil making soil unlikely to support certain native plants or more likely to favor non-native species) 10
- U. Unknown

Score

U

Documentation:
Identify ecosystem processes impacted (or if applicable, justify choosing answer A in the absence of impact information)
Applicable studies on impacts to ecosystem processes and system-wide parameters not known.
Sources of information:
Author's pers. comm.

1.2. Impact on Natural Community Structure

- A. No perceived impact; establishes in an existing layer without influencing its structure 0
- B. Influences structure in one layer (e.g., changes the density of one layer) 3
- C. Significant impact in at least one layer (e.g., creation of a new layer or elimination of an existing layer) 7
- D. Major alteration of structure (e.g., covers canopy, eradicating most or all layers below) 10
- U. Unknown

Score

3

Documentation:
Identify type of impact or alteration:
Has been observed to change the density of the herb layer.
Sources of information:
Author's pers. obs.

1.3. Impact on Natural Community Composition

- A. No perceived impact; causes no apparent change in native populations 0
- B. Influences community composition (e.g., reduces the number of individuals in one or more native species in the community) 3
- C. Significantly alters community composition (e.g., produces a significant reduction in the population size of one or more native species in the community) 7
- D. Causes major alteration in community composition (e.g., results in the extirpation of one or several native species, reducing biodiversity or change the community composition towards species exotic to the natural community) 10
- U. Unknown

Score

3

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

Documentation: Identify type of impact or alteration: Has been observed influencing community composition by reducing the number of native species in the community. Sources of information: Author's pers. obs.	
---	--

1.4. Impact on other species or species groups (cumulative impact of this species on the animals, fungi, microbes, and other organisms in the community it invades. Examples include reduction in nesting/foraging sites; reduction in habitat connectivity; injurious components such as spines, thorns, burrs, toxins; suppresses soil/sediment microflora; interferes with native pollinators and/or pollination of a native species; hybridizes with a native species; hosts a non-native disease which impacts a native species)

- | | |
|---|----|
| A. Negligible perceived impact | 0 |
| B. Minor impact | 3 |
| C. Moderate impact | 7 |
| D. Severe impact on other species or species groups | 10 |
| U. Unknown | |

Score

U

Documentation: Identify type of impact or alteration: Species is in <i>Cyperus</i> subg. <i>Pycnostachys</i> , which includes the native species, <i>Cyperus acuminatus</i> and <i>C. dentatus</i> . Hybridization between <i>C. difformis</i> and the native species of the subgenus are not known. Studies of the impacts on other species or species groups are not known Sources of information: Tucker et al., 2005; author's pers. comm.	
---	--

Total Possible	<table border="1" style="display: inline-table; width: 50px; height: 20px;"><tr><td style="text-align: center;">20</td></tr></table>	20
20		
Section One Total	<table border="1" style="display: inline-table; width: 50px; height: 20px;"><tr><td style="text-align: center;">6</td></tr></table>	6
6		

2. BIOLOGICAL CHARACTERISTICS AND DISPERSAL ABILITY

2.1. Mode and rate of reproduction

- | | |
|--|---|
| A. No reproduction by seeds or vegetative propagules (i.e. plant sterile with no sexual or asexual reproduction). | 0 |
| B. Limited reproduction (fewer than 10 viable seeds per plant AND no vegetative reproduction; if viability is not known, then maximum seed production is less than 100 seeds per plant and no vegetative reproduction) | 1 |
| C. Moderate reproduction (fewer than 100 viable seeds per plant - if viability is not known, then maximum seed production is less than 1000 seeds per plant - OR limited successful vegetative spread documented) | 2 |
| D. Abundant reproduction with vegetative asexual spread documented as one of the plants prime reproductive means OR more than 100 viable seeds per plant (if viability is not known, then maximum seed production reported to be greater than 1000 seeds per plant.) | 4 |
| U. Unknown | |

Score

4

Documentation: Describe key reproductive characteristics (including seeds per plant): Can produce hundreds of seeds with high viability. Sources of information: McIntyre, 1985; Tucker et al., 2005; Chauhan & Johnson, 2009; author's pers. obs.	
---	--

2.2. Innate potential for long-distance dispersal (e.g. bird dispersal, sticks to animal hair,

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

buoyant fruits, pappus for wind-dispersal)

- A. Does not occur (no long-distance dispersal mechanisms) 0
- B. Infrequent or inefficient long-distance dispersal (occurs occasionally despite lack of adaptations) 1
- C. Moderate opportunities for long-distance dispersal (adaptations exist for long-distance dispersal, but studies report that 95% of seeds land within 100 meters of the parent plant) 2
- D. Numerous opportunities for long-distance dispersal (adaptations exist for long-distance dispersal and evidence that many seeds disperse greater than 100 meters from the parent plant) 4
- U. Unknown

Score

4

Documentation:

Identify dispersal mechanisms:

Small achenes believed to be transported by birds. Lemaire (1970): "growing abundantly in a 'marsh' on the roof of the third floor and as scattered plants in very shallow water...on the sixth floor roof" of a building in Lincoln, Nebraska. Small achenes also could be readily transported by water.

Sources of information:

Lemaire, 1970; Tyndall, 1983

2.3. Potential to be spread by human activities (both directly and indirectly – possible mechanisms include: commercial sales, use as forage/revegetation, spread along highways, transport on boats, contaminated compost, land and vegetation management equipment such as mowers and excavators, etc.)

- A. Does not occur 0
- B. Low (human dispersal to new areas occurs almost exclusively by direct means and is infrequent or inefficient) 1
- C. Moderate (human dispersal to new areas occurs by direct and indirect means to a moderate extent) 2
- D. High (opportunities for human dispersal to new areas by direct and indirect means are numerous, frequent, and successful) 3
- U. Unknown

Score

2

Documentation:

Identify dispersal mechanisms:

Small achenes could be readily dispersed by mowing and farm equipment.

Sources of information:

Author's pers. obs.

2.4. Characteristics that increase competitive advantage, such as shade tolerance, ability to grow on infertile soils, perennial habit, fast growth, nitrogen fixation, allelopathy, etc.

- A. Possesses no characteristics that increase competitive advantage 0
- B. Possesses one characteristic that increases competitive advantage 3
- C. Possesses two or more characteristics that increase competitive advantage 6
- U. Unknown

Score

6

Documentation:

Evidence of competitive ability:

Plant exhibits fast growth and can grow on nutrient poor soils. Species is generally reported as an annual (Tucker et al., 2005), although some report (Holm et al., 1977; Imai & Murata, 1979) it to be an infrequent perennial.

Sources of information:

Holm et al., 1977; Imai & Murata, 1979; Tucker et al., 2005; author's pers. obs.

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

2.5. Growth vigor

- A. Does not form thickets or have a climbing or smothering growth habit 0
- B. Has climbing or smothering growth habit, forms a dense layer above shorter vegetation, forms dense thickets, or forms a dense floating mat in aquatic systems where it smothers other vegetation or organisms 2
- U. Unknown

Score

0

Documentation:

Describe growth form:
Does not form thickets or have a climbing or smothering growth habit.
Sources of information:
Author's pers. obs.

2.6. Germination/Regeneration

- A. Requires open soil or water and disturbance for seed germination, or regeneration from vegetative propagules. 0
- B. Can germinate/regenerate in vegetated areas but in a narrow range or in special conditions 2
- C. Can germinate/regenerate in existing vegetation in a wide range of conditions 3
- U. Unknown (No studies have been completed)

Score

2

Documentation:

Describe germination requirements:
Plants generally becomes established in open areas with limited existing vegetation. Subsequent generations of this annual species germinate in this existing vegetation. Chauhan & Johnson reported that seeds buried deeper than than 1 cm. do not germinate.
Sources of information:
Chauhan & Johnson, 2009; author's pers. obs.

2.7. Other species in the genus invasive in New York or elsewhere

- A. No 0
- B. Yes 3
- U. Unknown

Score

0

Documentation:

Species:
There are other non-native species of *Cyperus* (e.g., *Cyperus iria*) but none is tracked as invasive in New York or elsewhere. Brooklyn Botanic Garden, 2009; Weldy & Werier, 2009; U.S.D.A. NRCS, 2009.

Total Possible

25

Section Two Total

18

3. ECOLOGICAL AMPLITUDE AND DISTRIBUTION

3.1. Density of stands in natural areas in the northeastern USA and eastern Canada (use same definition as Gleason & Cronquist which is: "The part of the United States covered extends from the Atlantic Ocean west to the western boundaries of Minnesota, Iowa, northern Missouri, and southern Illinois, south to the southern boundaries of Virginia, Kentucky, and Illinois, and south to the Missouri River in Missouri. In Canada the area covered includes Nova Scotia, Prince Edward Island, New Brunswick, and parts of Quebec and Ontario lying south of the 47th parallel of latitude")

- A. No large stands (no areas greater than 1/4 acre or 1000 square meters) 0
- B. Large dense stands present in areas with numerous invasive species already present or 2

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

- disturbed landscapes
- C. Large dense stands present in areas with few other invasive species present (i.e. ability to invade relatively pristine natural areas) 4
- U. Unknown

Score

2

Documentation:

Identify reason for selection, or evidence of weedy history:
 Author has observed stands of 0.25 acre or more in Virginia. Some stands in New Jersey may approach 0.25 acre. In all cases, the landscape was disturbed with other invasive species being present.
 Sources of information:
 Author's pers. obs.

3.2. Number of habitats the species may invade

- A. Not known to invade any natural habitats given at A2.3 0
- B. Known to occur in one natural habitat given at A2.3 1
- C. Known to occur in two natural habitats given at A2.3 2
- D. Known to occur in three natural habitat given at A2.3 4
- E. Known to occur in four or more natural habitats given at A2.3 6
- U. Unknown

Score

6

Documentation:

Identify type of habitats where it occurs and degree/type of impacts:
 See A2.3.
 Sources of information:
 Tyndall, 1983; Tucker et al., 2005; Brooklyn Botanic Garden, 2009; author's pers. obs.

3.3. Role of disturbance in establishment

- A. Requires anthropogenic disturbances to establish. 0
- B. May occasionally establish in undisturbed areas but can readily establish in areas with natural or anthropogenic disturbances. 2
- C. Can establish independent of any known natural or anthropogenic disturbances. 4
- U. Unknown

Score

2

Documentation:

Identify type of disturbance:
 Always found in disturbed areas; not known to require anthropogenic disturbance.
 Sources of information:
 Tucker et al., 2005; author's pers. obs.

3.4. Climate in native range

- A. Native range does not include climates similar to New York 0
- B. Native range possibly includes climates similar to at least part of New York. 1
- C. Native range includes climates similar to those in New York 3
- U. Unknown

Score

1

Documentation:

Describe what part of the native range is similar in climate to New York:
 Native range includes climates similar to lower New York.
 Sources of information:
 Brooklyn Botanic Garden, 2009.

3.5. Current introduced distribution in the northeastern USA and eastern Canada (see question 3.1 for definition of geographic scope)

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

- A. Not known from the northeastern US and adjacent Canada 0
- B. Present as a non-native in one northeastern USA state and/or eastern Canadian province. 1
- C. Present as a non-native in 2 or 3 northeastern USA states and/or eastern Canadian provinces. 2
- D. Present as a non-native in 4–8 northeastern USA states and/or eastern Canadian provinces, and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 1 northeastern state or eastern Canadian province. 3
- E. Present as a non-native in >8 northeastern USA states and/or eastern Canadian provinces. and/or categorized as a problem weed (e.g., “Noxious” or “Invasive”) in 2 northeastern states or eastern Canadian provinces. 4
- U. Unknown

Score

3

Documentation:
 Identify states and provinces invaded:
 KY, PA, NJ, NY, VA.
 Sources of information: See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.
 Brooklyn Botanic Garden, 2009; U.S.D.A. NRCS, 2009.

3.6. Current introduced distribution of the species in natural areas in the eight New York State PRISMs (Partnerships for Regional Invasive Species Management)

- A. Present in none of the PRISMs 0
- B. Present in 1 PRISM 1
- C. Present in 2 PRISMs 2
- D. Present in 3 PRISMs 3
- E. Present in more than 3 PRISMs or on the Federal noxious weed lists 4
- U. Unknown

Score

1

Documentation:
 Describe distribution:
 See A1.1.
 Sources of information:
 Brooklyn Botanic Garden, 2009; Weldy & Werier.

Total Possible

25

 Section Three Total

15

4. DIFFICULTY OF CONTROL

4.1. Seed banks

- A. Seeds (or vegetative propagules) remain viable in soil for less than 1 year, or does not make viable seeds or persistent propagules. 0
- B. Seeds (or vegetative propagules) remain viable in soil for at least 1 to 10 years 2
- C. Seeds (or vegetative propagules) remain viable in soil for more than 10 years 3
- U. Unknown

Score

U

Documentation:
 Identify longevity of seed bank:
 Sources for length of seed viability for *Cyperus difformis* could not be located
 Sources of information:

**NEW YORK
NON-NATIVE PLANT INVASIVENESS RANKING FORM**

Author's pers. comm.

4.2. Vegetative regeneration

- | | | |
|----|---|---|
| A. | No regrowth following removal of aboveground growth | 0 |
| B. | Regrowth from ground-level meristems | 1 |
| C. | Regrowth from extensive underground system | 2 |
| D. | Any plant part is a viable propagule | 3 |
| U. | Unknown | |

Score 1

Documentation:

Describe vegetative response:
 Limited ground level growth has been observed after cutting.
 Sources of information:
 Author's pers. obs.

4.3. Level of effort required

- | | | |
|----|---|---|
| A. | Management is not required: e.g., species does not persist without repeated anthropogenic disturbance. | 0 |
| B. | Management is relatively easy and inexpensive: e.g. 10 or fewer person-hours of manual effort (pulling, cutting and/or digging) can eradicate a 1 acre infestation in 1 year (infestation averages 50% cover or 1 plant/100 ft ²). | 2 |
| C. | Management requires a major short-term investment: e.g. 100 or fewer person-hours/year of manual effort, or up to 10 person-hours/year using mechanical equipment (chain saws, mowers, etc.) for 2-5 years to suppress a 1 acre infestation. Eradication is difficult, but possible (infestation as above). | 3 |
| D. | Management requires a major investment: e.g. more than 100 person-hours/year of manual effort, or more than 10 person hours/year using mechanical equipment, or the use of herbicide, grazing animals, fire, etc. for more than 5 years to suppress a 1 acre infestation. Eradication may be impossible (infestation as above). | 4 |
| U. | Unknown | |

Score 0

Documentation:

Identify types of control methods and time-term required:
 While the species requires control in other areas (e.g., areas where rice is cultivated) management is not currently required in New York.
 Sources of information:
 Author's pers comm.

Total Possible	7
Section Four Total	1

Total for 4 sections Possible	77
Total for 4 sections	40

C. STATUS OF CULTIVARS AND HYBRIDS:

At the present time (May 2008) there is no protocol or criteria for assessing the invasiveness of cultivars independent of the species to which they belong. Such a protocol is needed, and individuals with the appropriate expertise should address this issue in the future. Such a protocol will likely require data on cultivar fertility and identification in both experimental and natural settings.

Hybrids (crosses between different parent species) should be assessed individually and separately from the parent species wherever taxonomically possible, since their invasiveness may differ from that of the parent species. An exception should be made if the taxonomy of the species and hybrids are uncertain,

NEW YORK NON-NATIVE PLANT INVASIVENESS RANKING FORM

and species and hybrids can not be clearly distinguished in the field. In such cases it is not feasible to distinguish species and hybrids, and they can only be assessed as a single unit.

Some cultivars of the species known to be available:

References for species assessment:

Brooklyn Botanic Garden. 2008. AILANTHUS database. [Accessed on July 20, 2009.]

Chauhan, B. S. and D. E. Johnson. Ecological studies on *Cyperus difformis*, *Cyperus iria*, and *Fimbristylis miliacea*: three troublesome annual sedge weeds of rice. *Annals of Applied Biology* 155(1): 103-112.

Holm, L. G., D. L. Plucknett, J. V. Pancho, and J. P. Herberger. 1977. *The world's worst weeds; distribution and biology*. University of Georgia Press, Athens. 712 pp.

Imai, K. and Y. Murata. 1979. Changes in apparent photosynthesis, CO₂ compensation point and dark respiration of leaves of some Poaceae and Cyperaceae species with senescence. *Plant Cell Physiology* 20: 1653-1658.

Lemaire, R. J. 1970. Recent plant records for Nebraska. *Rhodora* 72: 283-284.

McIntyre, S. 1985. Seed reserves in temperate Australian rice field following rotation and continuous cropping. *Journal of Applied Ecology* 22: 875-884.

Tucker, G. C., B. G. Marcks, and J. R. Carter. 2002. *Cyperus* Pp. 141-191 in in Ball et al. (eds.), *Flora of North America*. Vol. 23. Oxford University Press, New York. 608 pp.

Tyndall, R. W. 1983. Distribution of *Cyperus difformis* L. (Cyperaceae) in the southeastern United States. *Castanea* 48(4): 277-280.

United States Department of Agriculture, National Resources Conservation Service. 2008. The PLANTS Database. National Plant Data Center, Baton Rouge, Louisiana. [Accessed on July 20, 2009.]

Weldy, T. and D. Werier. 2005. *New York Flora Atlas*. [S.M. Landry, K.N. Campbell, and L.D. Mabe (original application development), Florida Center for Community Design and Research. University of South Florida]. New York Flora Association, Albany, New York. [Accessed on July 20, 2009].

Zimdahl, R. L. K. Moody, R. T. Lubigan, E. M. castin. 1988. Patterns of weed emergence in tropical soil. *Weed Science* 36: 603-608.

Citation: This NY ranking form may be cited as: Jordan, M.J., G. Moore and T.W. Weldy. 2008. Invasiveness ranking system for non-native plants of New York. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, NY. Note that the order of authorship is alphabetical; all three authors contributed substantially to the development of this protocol.

Acknowledgments: The NY form incorporates components and approaches used in several other systems, cited in the references below. Valuable contributions by members of the Long Island Invasive Species Management Area's Scientific Review Committee were incorporated in revisions of this form. Original members of the LIISMA SRC included representatives of the Brooklyn Botanic Garden; The Nature Conservancy; New York Natural Heritage

NEW YORK

NON-NATIVE PLANT INVASIVENESS RANKING FORM

Program, New York Sea Grant; New York State Office of Parks, Recreation and Historic Preservation; National Park Service; Brookhaven National Laboratory; New York State Department of Environmental Conservation Region 1; Cornell Cooperative Extension of Suffolk/Nassau Counties; Long Island Nursery and Landscape Association; Long Island Farm Bureau; SUNY Farmingdale Ornamental Horticulture Department; Queens College Biology Department; Long Island Botanical Society; Long Island Weed Information Management System database manager; Suffolk County Department of Parks, Recreation and Conservation; Nassau County Department of Parks, Recreation and Museums; Suffolk County Soil & Water Conservation District.

References for ranking form:

- Carlson, Matthew L., Irina V. Lapina, Michael Shephard, Jeffery S. Conn, Roseann Densmore, Page Spencer, Jeff Heys, Julie Riley, Jamie Nielsen. 2008. Invasiveness ranking system for non-native plants of Alaska. Technical Paper R10-TPXX, USDA Forest Service, Alaska Region, Anchorage, AK XX9. Alaska Weed Ranking Project may be viewed at: http://akweeds.uaa.alaska.edu/akweeds_ranking_page.htm.
- Heffernan, K.E., P.P. Coulling, J.F. Townsend, and C.J. Hutto. 2001. Ranking Invasive Exotic Plant Species in Virginia. Natural Heritage Technical Report 01-13. Virginia Dept. of Conservation and Recreation, Division of Natural Heritage, Richmond, Virginia. 27 pp. plus appendices (total 149 p.).
- Morse, L.E., J.M. Randall, N. Benton, R. Hiebert, and S. Lu. 2004. An Invasive Species Assessment Protocol: Evaluating Non-Native Plants for Their Impact on Biodiversity. Version 1. NatureServe, Arlington, Virginia. <http://www.natureserve.org/getData/plantData.jsp>
- Randall, J.M., L.E. Morse, N. Benton, R. Hiebert, S. Lu, and T. Killeffer. 2008. The Invasive Species Assessment Protocol: A Tool for Creating Regional and National Lists of Invasive Nonnative Plants that Negatively Impact Biodiversity. *Invasive Plant Science and Management* 1:36–49
- Warner, Peter J., Carla C. Bossard, Matthew L. Brooks, Joseph M. DiTomaso, John A. Hall, Ann M. Howald, Douglas W. Johnson, John M. Randall, Cynthia L. Roye, Maria M. Ryan, and Alison E. Stanton. 2003. Criteria for Categorizing Invasive Non-Native Plants that Threaten Wildlands. Available online at www.caleppc.org and www.swvma.org. California Exotic Pest Plant Council and Southwest Vegetation Management Association. 24 pp.
- Williams, P. A., and M. Newfield. 2002. A weed risk assessment system for new conservation weeds in New Zealand. *Science for Conservation* 209. New Zealand Department of Conservation. 1-23 pp.